

Relationship between Anemia and Sore mouth

Original Article

Donia Sadri¹, Assadollah Moosavi², Emelia Mohaghegh³

¹ Associate Professor, Department of Oral & Maxillofacial Pathology, Dental Branch, Islamic Azad University, Tehran, Iran.

² Associate Professor, Department of Hematology & Oncology, Tehran University of Medical Sciences, Shariati Hospital, Tehran, Iran.

³ Dentist

Received: Sep 4, 2011

Accepted: Jan 18, 2012

Corresponding Author:

Donia Sadri, DDS, Msc

Address: Department of Oral & Maxillofacial Pathology, Dental Branch, Islamic Azad University, Tehran, Iran.

Telephone: +98-21- 22763449

Fax: +98-21-22763432

E-mail: Donia1351@yahoo.com

Abstract

Introduction: According to the prevalence of sore mouth, its outcome and possible role of anemia in its etiology and some contradictions in this theory, our study was performed to evaluate the role of iron deficiency anemia as a etiological factor for sore mouth.

Materials and Methods: This historical cohort study was performed on 40 cases, including 20 anemic patients as the experimental group and 20 healthy individuals as the control group with similarity in socio economical, sex and age factors. Oral examination was carried out in two groups to evaluate soremouth presentations including atrophic glossitis, recurrent oral ulceration, pale oral mucosa, tongue burning sensation, glossodynia and angular cheilitis. Data was analyzed by chi-square test. Attributable risk (AR) and related risk (RR) were calculated.

Results: Out of 40 studied cases, sore mouth manifestations were found in (45%) of individuals in case and (5%) control groups ($p < 0.005$). Regarding the anemic patients, RR and AR of sore mouth were 9 and 40% respectively. The most prevalent manifestations of sore mouth were atrophic glossitis, recurrent oral ulceration, pale oral mucosa, tongue burning sensation, glossodynia and angular cheilitis.

Conclusion: According to this study, it seems iron deficiency anemia has a significant role in sore mouth incidence. Evaluation of anemia treatment effect on sore mouth elimination is recommended.

Key words: •Anemia •Oral ulcer •Cheilitis •Mucositis

Introduction

Sore mouth is a group of oral lesions including atrophic glossitis, prickly sensation of tongue, pale mucosa, recurrent aphthous ulcer, angular cheilitis, erythematous mucositis and glossodynia.⁽¹⁾

Blood deficiency and malnutrition are the two major proposed etiological factors for sore mouth.⁽²⁾

In the recent years, iron deficiency anemia has been prevalent in our country, World Health Organization(WHO) reported that 58 percent of woman in undeveloped countries are suffering from anemia.^(3,4) Anemia with blood hemoglobin decrease leading to reduced tissue oxygenation can result in undesirable effects.⁽⁵⁾

It has been argued that the treatment with ferrous supplement in iron deficiency anemia of the patients with atrophic glossitis symptoms, can reduce the oral lesions.^(6,7)

Dali-Youcef et al found out that the patients with megaloblastic anemia are more susceptible to osteomyelitis and atrophic glossitis than the healthy ones.⁽⁸⁾

However, results of the most influential research about the relationship between oral mucosa lesions and anemia were inconclusive and other studies have also found conflicting results.⁽⁹⁻¹¹⁾

Materials and Methods

This historical cohort study was conducted on 20 consecutive patients with anemia as the experimental group of 9 males, 11 females, aged (41.7±12.6) and 20 healthy ones as the control group of 8 males, and 12 females, aged(40.5±15.08) that was matched with the experimental group.

All participants referred to Shariati hospital Tehran-Iran from January 2010 to May 2010.

The ethical committee of the university approved the study (No:P/165/d). Anemia was confirmed by paraclinical tests, consist-

ing of Fe, Ferritin, RDW-CV, HCT, MCH, MCV, Hb, RBC, TIBC. All participants were evaluated by oral pathologist for diagnosis of sore mouth sign and symptoms (atrophic glossitis, prickly sensation of tongue, pale mucosa, recurrent aphthous ulcer,angular cheilitis, erythematous mucositis and glossodynia.⁽¹²⁾

The data were analyzed using Chi-square test in SPSS version 16. P value of 0.05 was considered indicative of a statistically significant difference. Attributable risk (AR) and related risk (RR) were calculated, too.

Results

In this study, we evaluated sore mouth signs and symptoms in 20 anemic and 20 healthy participants. Table 1 indicates the correlations between anemia and sore mouth (P<0.05).

Related risk (RR) for incidence of sore mouth in anemic patients was 9 times more than control group (RR= 9).

Attributable risk (AR) of sore mouth incidence in anemic patients was 40 percent. (AR= 40%)




Figure 1. Illustrates the signs and symptoms of sore mouth in 20 anemic patients

- 1- AG= Atrophic glossitis
- 2- ROU=Recurrent oral ulceration
- 3- POM=Pale oral mucosa
- 4- TBS=Tongue burning sensation
- 5- GD=glossodynia
- 6- AC=Angular cheilitis

Table 1. Prevalence of sore mouth in experimental and control groups

Groups	Sore mouth Not found No (%)	Found No (%)	Total	P-value
Control	19 (95)	1(5)	20	<0.05
Experimental	11(55)	9 (45)	20	

Discussion

According to the results of the study, anemia had a positive correlation with sore mouth. The results are in agreement with those of Lu Sy et al.⁽¹²⁾ One possible explanation for this relationship is that anemia with hemoglobin decrease leads to the decreasing of oxygenation to oral soft tissues that leads to sore mouth.^(13,14)

However, the results are in disagreement with those of Vucicevic-Boras et al's study, in that, they found no correlation between anemia and sore mouth.⁽¹⁴⁾

The possible justification for this discrepancy is the difference in the accuracy of lab tests and neglecting the confounding factors.

In this study; out of 20 anemic patients, 9 cases (45%) have signs and symptoms of sore mouth. The results are in agreement with those of Petavy-Catala et al.⁽¹⁵⁾

In our study, 20% of the experimental group had recurrent aphthous ulcer and only 5% of the control group had this lesion. Although this relationship is not statistically significant, this result has also been observed in the previous studies.⁽¹⁶⁻²⁰⁾

In this study, 25% of the experimental group had glossitis with a different degree of papillary atrophy and nobody in the control group had this lesion. Despite the fact that this relationship was not statistically significant, our results support those of Graells's study.⁽²⁰⁾

In our study, the prevalence of glossodynia and glossitis in the experimental group was 15% and 10% respectively. The present findings show more prevalence of these lesions to the other studies.^(21,22) A possible explanation for this finding is high proliferation rate of lingual mucosa and demanding of acid folic and ferrous for cell proliferation. Decreasing of these micronutrients elements can also facilitate candidal proliferation and candidiasis incidence that lead to glossitis and glossodynia.

Conclusion

According to this study, it seems iron deficiency anemia has a significant role in sore mouth incidence. Evaluation of anemia treatment effect on sore mouth elimination is recommended.

References

1. Wood NK, Goaz PW. Differential diagnosis of oral and maxillofacial lesions. 7th ed. St Louis Missouri, Mosby Inc; 2000, pp:583-9.
2. Neville BW, Damm DD, Allen CM, Bouquot JE. Oral and maxillofacial pathology. 3rd ed. Philadelphia: WB Saunders Company; 2009. pp: 341-5.
3. Ludwig H, Strasser K. Symptomatology of anemia. *Semin Oncol* 2001; 28: 7-14.
4. Piskin S, Sayan C, Durukan N, Senson L M. Serum iron, ferritin: Folic acid and cobalamine levels in recurrent aphthous stomatitis. *J Eur Acad Dermatol Venerol* 2002; 16: 66-67.
5. Porter SR, Scully C, Flint S. Hematologic status in recurrent aphthous stomatitis compared with other oral disease. *Oral Surg Oral Med Oral Pathol* 1988; 66: 41-4.
6. Vidal-alaballa J, Butler CC, Cannings-John R, et al. Oral vitamin B12 versus intramuscular vitamin B12 for vitamin B12 deficiency. *Cochrane Database Syst Rev* 2005; (3): CD004655.
7. Tyldesley WR. Stomatitis and recurrent oral ulceration: is a full blood screen necessary. *Br J Oral Surg* 1983; 21: 27-30.
8. Dali-Youcef N, Andres E. An update on cobalamin deficiency in adults. *QJM* 2009 ; 102(1): 17-28.
9. Longo DI, Fauci A, Kasper D, et al. *Harrisons principles of Internal Medicine*, 18th ed, New York, McGraw-Hill Inc; 2011. pp: 660-74.
10. Oh RC, Brown DL. Vitamin B12 deficiency. *Am Fam Physician* 2003; 67(5): 979-86.
11. Enwald CV, Darinka PJ, Swartz C, et al. Iron status in atrophic glossitis: a pilot study. *Wis Med J*. 1993; 92(10): 570-3.

Anemia and Sore mouth

12. Lu Sy, Wu HC. Initial diagnosis of anemia from sore mouth and improved classification of anemia by MCV, RDW in 30 patients. *Oral med oral pathol oral radiolendod* 2004; 98: 679-85.
13. Kobkan T, PorpanY, Viliawan A. Folate and Vit B12 levels in patients with oral Lichen planus stomatitis or glossitis. *South east Asian J* 2001; 32: 643-647.
14. Vucicevic-Boras V, Topic B, Cekic-Armbasin A, et al. Lack of association between burning mouth and hematinic deficiencies. *Eur J Med Res* , 2001; 28: 6(9):409-12.
15. Petavy- Catala C, Fonts V, Granet N. Clinical manifestation of the mouth revealing vit B12 deficiency before the onset of anemia. *Ann Dermatol Venerol* 2003; 130: 191-4.
16. Kozlak St, Walsh SJ, Lalla RV. Reduced dietary take of vit B12 and folate in patient with aphtous stomatitis. *J oral Pathol Med* 2010; 39(5):420-3.
17. PierroVS, Maia LC, Primo LF, Soares FD. Case report: the importance of oral manifestations in diagnosing iron deficiency in childhood. *Eur J Pediatric Dent* 2004; 5(2): 115-8.
18. Gao J, Chen L, Zhou J, Peng J. A case control study on etiological factors involved in patients with burning mouth syndrome. *J oral pathol med*2009; 38: 24-28.
19. DrinkaPj, LangerE, ScottL, MorrowF. Laboratory measurement of nutritional status as correlates of atrophic glossitis. *J Gen Intern Med* 1991; 6: 37-140.
20. Jordi G, Rosa MO, Cristiana M. Glossitis with linear lesions: an early sign of vit B12 deficiency *JAM Acad Dermatol* 2009; 6: 489-500.
21. Pontes HA, Neto NC, Ferreira KB, et al. Oral manifestation of vit B12 deficiency: a case report. *J Can Den Assoc* 2009; 75(7): 533-7.
22. Adeyemo TA, Adeyemo WL, Adediran A, et al. Orofacial manifestation of hematological disorders: Anemia and hemostatic disorders. *Indian J Dent Res*2011;22(3):454-61.