

Research Paper: Comparison of intra oral manifestations of pregnant and non-pregnant women: an observational case-control study



Zahra Elmi Rankouhi¹, Seyed Javad Kia¹, Dina Maleki^{2*}, Mohsen Masoumi³

¹Dental Sciences Research Center, Department of Oral and Maxillofacial Medicine, School of Dentistry, Gilan University of Medical Sciences, Rasht, Iran

²Dentistry Student, Student Research Committee, School of Dentistry, Gilan University of Medical Sciences, Rasht, Iran

³Dentist

Use your device to scan
and read the article online



Citation: Rankouhi Elmi Z, Kia SJ, Maleki D, Masoumi M. Comparison of intra oral manifestations of pregnant and non-pregnant women: an observational case-control study. Journal of Dentomaxillofacial Radiology, Pathology and Surgery. 2019; 8(3):34-38. <http://dx.doi.org/10.32598/3dj.7.4.145>

<http://3dj.gums.ac.ir>



ABSTRACT

Introduction: Hormonal changes during pregnancy are along with oral changes such as pregnancy gingivitis, halitosis, pregnancy tumor and teeth erosion. The aim of the current study is to evaluate the prevalence of these oral changes in pregnant and non-pregnant women.

Materials and Methods: In this observational, analytic, case-control study, 124 pregnant women and 124 non-pregnant women referred to Al-zahra hospital, Rasht, Iran, were examined. Age, education, number of pregnancy and pregnant status of patients were recorded. Also, gingival index, plaque index, halitosis, pregnancy tumor, erosion and geographic tongue were assessed and recorded. Statistical analysis was performed with SPSS 22. Data were analyzed using the Chi square and Mann-Whitney tests with $P < 0.05$ considered significant.

Results: No statistical relation was found between age and pregnancy status. ($P=0.085$) Also, the relation of education and pregnancy status was not statistically significant. ($P=0.49$) Plaque index, gingival index and halitosis were significantly different between pregnant and non-pregnant women. ($P=0.018$, $P=0.001$ and $P=0.0001$ respectively) However, pregnancy tumor, erosion and geographic tongue were not significantly different based on pregnancy status. ($P=0.65$, $P=0.758$ and $P=0.23$)

Conclusion: During pregnancy, occurrence of gingival inflammation and halitosis increases based on the current study. It should be noted that better oral hygiene is of benefits for pregnant patients, offering them comfort, function and aesthetics.

Article info:

Received: 2019/08/23

Accepted: 2019/09/01

Keywords:

Pregnancy
Halitosis
Tooth Attrition
Tooth Wear
Periodontal Diseases

* Corresponding Author:

Dinya Maleki .

Address: Department of Oral and Maxillofacial Surgery, Guilan University of Medical Sciences.

Tel: +989128998663

E-mail: dinamaleki74@gmail.com

Introduction

Oral health assess during pregnancy is critical since poor oral hygiene can lead to adverse problems such as low birthweight, preterm birth, and preeclampsia. (1-2) Also, hormonal changes during pregnancy may result in oral mucosal changes such as pregnancy gingivitis, halitosis, pregnancy tumor, teeth erosion, geographic tongue, chloasma, facial telangiectasia, sialorrhea, tooth mobility and oral apathies. (2)

Pregnancy gingivitis is the gingival inflammation which results from dental plaque accumulation and is exacerbated by pregnancy hormones. (3) 50 to 100% of pregnant women experience pregnancy gingivitis and almost 100% of pregnant women need periodontal treatment during their pregnancy. (1,4) Halitosis during pregnancy is the result of hormonal change and pregnancy gingivitis. Also, halitosis in pregnant women, is associated to periodontal disease, caries, xerostomia, vomiting, gastric reflux and gestational diabetes. (5) Pregnancy tumor is an exaggerated inflammatory response to an irritation which begin during the first trimester, and worsens as the pregnancy progresses up through the seventh month of pregnancy. (4-6) The prevalence of pregnancy tumor has been reported to vary between 0.5-100% in different studies. (1,4-5) Nausea and vomiting in pregnancy may lead to extensive erosions specially affecting the palatal surfaces of the maxillary incisors and canines. (2,5) Following the erosion of teeth, hyper sensitivity is a common finding as a result of dentin exposure. (3,5) Geographic tongue occurs because of the atrophy of the filiform papillae. Genetic factors, hormonal changes and oral contraceptive pills, pregnancy and diabetes mellitus are possible risk factors of geographic tongue. (7,8)

The aim of the current study was to evaluate the prevalence of potential intra-oral manifestations in pregnant and non-pregnant women.

Methods and materials

In this observational, analytic, case-control study, 124 pregnant women and 124

non-pregnant women referred to Al-zahra hospital, Rasht, Guilan, Iran, were examined.

Pregnant and non-pregnant women with no previous history or presence of systemic disease, aged between 20–45 years were included in the study as case group and control group, respectively. Patients who use contraceptive drugs, tobacco and alcohol, patients suffering from ovarian cyst or fibroma, and patients who are in their first 6 months after giving birth were excluded from the study. Written informed consents were provided from the subjects.

Age, education and pregnant status of patients were recorded. Age was documented as under 30 and over 30 years old. Education was determined as “up to diploma”, “bachelor” and “more than bachelor”. Pregnancy status was considered as being pregnant or not being pregnant.

Also, presence of gingival inflammation, halitosis, pregnancy tumor, erosion and geographic tongue were assessed and recorded.

Gingival inflammation was evaluated using O’Leary plaque index (9) and Löe and Silness gingival index (10). To determine O’Leary index, a disclosing agent was used and the stained surfaces of each tooth was recorded. O’Leary index was reported as the ratio of the total number of stained surfaces to the total number of existing surfaces. To assess Leo and Silness gingival index, a blunt explorer was used. Semi-erupted teeth, remaining root and teeth with peri apical lesion were not evaluated. Gingival index was recorded for each surface of a tooth as 0 (No inflammation), 1 (Mild inflammation, no bleeding on probing), 2 (Moderate inflammation, bleeding on probing) and 3 (severe inflammation, spontaneous bleeding). The sum of scores divided by the number of total surfaces was calculated. If the score was from 0.1 to 1, it would be described as mild gingivitis. If the score was from 1.1 to 2 and from 2.1 to 3, it would be described as moderate gingivitis and severe gingivitis, respectively.

To assess halitosis, Organoleptic Index defined by Rosenberg, was used. (11) The index is recorded as 0 to 5. After accurate examination of soft tissue, palatal surface of maxillary incisor teeth

and tongue, existence of pregnancy tumor, tooth erosion and geographic tongue were recorded.

Statistical analysis was performed with SPSS 22. Data were analyzed using the Chi square and Mann-Whitney tests with $P < 0.05$ considered significant.

Results

The mean age of samples was 30.95 ± 6.34 , the mean age of pregnant women and non-pregnant women was 30.25 ± 5.68 and 31.64 ± 6.89 , respectively. Data distribution of patients is presented in table 1.

Table 1: Data distribution of demographic information of patients.

		Pregnancy status		Total
		Pregnant	Non-Pregnant	
Age	Under 30 years old	50% (62)	48.4% (60)	49.2% (122)
	Over 30 years old	50% (62)	51.6% (64)	50.8% (126)
Education	Up to diploma	78.2% (97)	77.4% (96)	77.8% (193)
	Bachelor	21% (26)	19.4% (24)	20.2% (50)
	More than bachelor	0.8% (1)	3.2% (4)	2% (5)

No statistical relation was found between age and pregnancy status. ($P=0.085$) Also, the relation of education and pregnancy status was not statistically significant. ($P=0.49$)

Plaque index, gingival index and halitosis were significantly different between pregnant and non-pregnant women. ($P=0.018$, $P=0.001$ and $P=0.0001$ respectively) As such, the above-mentioned parameters were significantly higher in pregnant women compared to non-pregnant women. However, pregnancy tumor, erosion and geographic tongue were not significantly different based on pregnancy status. ($P=0.65$, $P= 0.758$ and $P=0.23$) (table 2)

Table 2: Data distribution of plaque index, gingival index, erosion, pregnancy tumor, geographic tongue, halitosis.

		Pregnancy status		Total
		Pregnant	Non-Pregnant	
O'Leary Plaque index	Less than 50%	0 (0)	1.6% (2)	0.8% (2)
	51-70%	0 (0)	8.1% (10)	4% (10)
	71-90%	41.1% (51)	41.1% (51)	41.1% (102)
	91-100%	58.9% (73)	49.2% (61)	54% (134)
Gingival index	Score 0	0 (0)	0 (0)	0 (0)
	Score 1	31.5% (39)	52.4% (65)	41.9% (104)
	Score 2	66.1% (82)	47.6 % (59)	56.9% (141)
	Score 3	2.4% (3)	0 (0)	1.2% (3)
Erosion	With erosion	4.8% (6)	4 % (5)	4.4% (11)
	Without erosion	95.2% (118)	96% (119)	95.6% (237)
Pregnancy tumor	With pregnancy tumor	2.4% (3)	1.6% (2)	2% (5)
	Without pregnancy tumor	97.6% (121)	98.4 % (122)	98% (243)
Geographic tongue	With geographic tongue	6.5% (8)	3.2% (4)	4.8% (12)
	Without geographic tongue	93.5% (116)	96.8% (120)	95.2% (236)
Halitosis	Score 0	18.5% (23)	54% (67)	36.3% (90)
	Score 1	15.3% (19)	17.7% (22)	16.5% (41)
	Score 2	31.5% (39)	16.9% (21)	24.2% (60)
	Score 3	31.5% (39)	11.3% (14)	21.4% (53)
	Score 4	3.2% (4)	0 (0)	1.6% (4)
	Score 5	0 (0)	0 (0)	0 (0)

Discussion

Estrogen and progesterone increase during pregnancy. Estrogen blocks gingival tissue turnover and its ability to produce collagen which results to an increased response to plaque accumulation. (1) On the other hand, progesterone prevents mitogen activation, cytotoxic

T cell generation and interleukin-6 production which make gingiva less resistant to inflammation. In conclusion, estrogen and progesterone make gingiva of pregnant women more susceptible to periodontal breakdown. (2)

In the current study, it was found that periodontal condition of pregnant women was worse than non-pregnant women which is reflected by measuring and comparing gingival index and plaque index. Lopez et al., Jain et al., Pentapati et al., Machuca et al., Soory, Mascarenhas et al., Chung et al. and Gonzales et al. reported occurrence of gingivitis to be higher in pregnant women compared to non-pregnant women. (12-19) Cohen et al., Miyazaki et al. and Maybodi et al. found that gingivitis gradually worsens from first semester to third semester in pregnant women. (20-22) However, Tilakaratne et al. found no significant difference between pregnant and non-pregnant women in term of plaque index values. (23) Also, Figuero et al. found no relation between gingival index and pregnancy. (24)

Physiologic and hormonal changes during pregnancy can also decrease saliva PH and increase bacterial growth, consequently leading to halitosis and higher risk of caries. (1,7) In the current study and Fujiwara et al. study, occurrence of halitosis was significantly higher in pregnant women compared to non-pregnant women. (25) Kia et al. reported that 37.7% of pregnant women experience halitosis during pregnancy. (8)

Pregnancy stimulates local production of angiogenic factors (vascular endothelial growth factor) in oral mucosal leading to the formation of pregnancy tumor. (26-27) Cardoso et al. observed an association between pregnancy tumor and pregnancy. (28) While, the current study found no statistical relation between pregnancy tumor and pregnancy, and reported that only 2.4% of pregnant patients had pregnancy tumor. Also, Chamani et al., Pirie et al. and Kia et al. found that respectively, 4.2%, 5% and 1% of pregnant patients had pregnancy tumor. (8, 29-30)

The current study reported dental erosion to have no significant relation with pregnancy status while AL-Sultani stated that erosion was significantly associated with pregnancy special-

ly in the third semester. (31) Kia et al. found that only 1.3% of patients had dental erosion. (8)

In accordance with the current study, Sarifakioglu et al. revealed no significant difference between pregnant and non-pregnant women in term of geographic tongue. (32) However, as opposed, Dvaz-Guzman et al. reported occurrence of geographic tongue to be more common in pregnant women. (7)

conclusion

During pregnancy, occurrence of gingival inflammation and halitosis increases based on the current study. It should be noted that better oral hygiene is of benefits for pregnant patients, offering them comfort, function and aesthetics.

References

1. Silva de Araujo Figueiredo C, Gonçalves Carvalho Rosalem C, Costa Cantanhede AL, Abreu Fonseca Thomaz ÉB, Fontoura Nogueira da Cruz MC. Systemic alterations and their oral manifestations in pregnant women. *Journal of Obstetrics and Gynaecology Research*. 2017; 43(1):16-22. <https://doi.org/10.1111/jog.13150>
2. Khanna S, Shalini M. Pregnancy and oral health: forgotten territory revisited. *The Journal of Obstetrics and Gynecology of India*. 2010; 60(2):123-7. <https://doi.org/10.1007/s13224-010-0017-0>
3. Usin MM, Tabares SM, Parodi RJ, Sembaj A. Periodontal conditions during the pregnancy associated with periodontal pathogens. *Journal of investigative and clinical dentistry*. 2013; 4(1):54-9. <https://doi.org/10.1111/j.2041-1626.2012.00137.x>
4. Nair V, Giri PK, Tirthankar D, Rudra A. Oral pregnancy tumor in pregnancy: a predicament. *IMJ*. 2016; 23(1):64-5.
5. Skouteris CA, editor. *Dental Management of the Pregnant Patient*. Wiley Blackwell; 2018. <https://doi.org/10.1002/9781119286592>
6. Barzegar M, Pouyafard A, Tabatabaei SH, Navabazam A, Ajami H, Chochedri M. Pregnancy tumor and facial port-wine stain: A case report. *Journal of Cranio-maxillofacial Research*. 2018 ; 22(1):50-3.
7. Diaz-Guzman LM, Castellanos-Suárez JL. Lesions of the oral mucosa and periodontal disease behavior in pregnant patients. *Medicina oral, patologia oral y cirugia bucal*. 2004;9(5):434-7.
8. Kia SJ, Vadiati Saberi B, Moonesan MR, Madani AR. Frequency of intra and extra oral manifestations of pregnant women in Rasht (2013). *The Journal of Qazvin*

- University of Medical Sciences. 2017; 20(6):12-8.
9. O'Leary TJ, Drake RB, Naylor JE. The plaque control record. *Journal of periodontology*. 1972; 43(1):38-. <https://doi.org/10.1902/jop.1972.43.1.38>
 10. Silness J, L oe H. Periodontal disease in pregnancy: Iii. Response to local treatment. *Acta odontologica scandinavica*. 1966; 24(6):747-59. <https://doi.org/10.3109/00016356609028739>
 11. Rosenberg M, Septon I, Eli I, Bar-Ness R, Gelernter I, Brenner S, Gabbay J. Halitosis measurement by an industrial sulphide monitor. *Journal of periodontology*. 1991; 62(8):487-9. <https://doi.org/10.1902/jop.1991.62.8.487>
 12. L opez LM, Guerra ME. Caries experience and periodontal status during pregnancy in a group of pregnant women with HIV+ infections from puerto rico. *Journal of AIDS & clinical research*. 2015; 6(3):45-51. <https://doi.org/10.4172/2155-6113.1000434>
 13. Jain K, Kaur H. Prevalence of oral lesions and measurement of salivary pH in the different trimesters of pregnancy. *Singapore medical journal*. 2015; 56(1):53-8. <https://doi.org/10.11622/smedj.2015010>
 14. Pentapati KC, Acharya S, Bhat M, Krishna Rao SV, Singh S. Oral health among women-a cross sectional study from South India. *International Journal of Dental Sciences and Research*. 2015;3(1):17-9.
 15. Machuca G, Khoshfeiz O, Lacalle JR, Machuca C, Bull on P. The influence of general health and socio-cultural variables on the periodontal condition of pregnant women. *Journal of periodontology*. 1999; 70(7):779-85. <https://doi.org/10.1902/jop.1999.70.7.779>
 16. Soory M. Hormonal factors in periodontal disease. *Dent Update*. 2000; 27(1):380-383. <https://doi.org/10.12968/denu.2000.27.8.380>
 17. Mascarenhas P, Gapski R, Al-Shammari K, Wang HL. Influence of sex hormones on the periodontium. *Journal of clinical periodontology*. 2003; 30(8):671-81. <https://doi.org/10.1034/j.1600-051X.2003.00055.x>
 18. Chung LH, Gregorich SE, Armitage GC, Gonzalez-Vargas J, Adams SH. Sociodemographic disparities and behavioral factors in clinical oral health status during pregnancy. *Community dentistry and oral epidemiology*. 2014;42(2):151-9. <https://doi.org/10.1111/cdoe.12075>
 19. Gonz alez-Jaranay M, T ellez L, Roa-L opez A, G omez-Moreno G, Moreu G. Periodontal status during pregnancy and postpartum. *PloS one*. 2017;12(5):45-50. <https://doi.org/10.1371/journal.pone.0178234>
 20. Cohen DW, Friedman L, Shapiro J, Kyle GC. A longitudinal investigation of the periodontal changes during pregnancy. *Journal of Periodontology-Periodontics*. 1969;40(10):563-70. <https://doi.org/10.1902/jop.1969.40.10.563>
 21. Miyazaki H, Yamashita Y, Shirahama R, Goto-Kimura K, Shimada N, Sogame A, Takehara T. Periodontal condition of pregnant women assessed by CPITN. *Journal of clinical periodontology*. 1991;18(10):751-4. <https://doi.org/10.1111/j.1600-051X.1991.tb00067.x>
 22. Maybodi FR, Haerian-Ardakani A, Vaziri F, Khabbazian A, Mohammadi-Asl S. CPITN changes during pregnancy and maternal demographic factors 'impact on periodontal health. *Iranian journal of reproductive medicine*. 2015;13(2):107.
 23. Tilakaratne A, Soory M, Ranasinghe AW, Corea SM, Ekanayake SL, De Silva M. Periodontal disease status during pregnancy and 3 months post-partum, in a rural population of Sri-Lankan women. *Journal of clinical periodontology*. 2000; 27(10):787-92. <https://doi.org/10.1034/j.1600-051x.2000.027010787.x>
 24. Figuero E, Carrillo-de-Albornoz A, Mart ın C, Tob as A, Herrera D. Effect of pregnancy on gingival inflammation in systemically healthy women: a systematic review. *Journal of clinical periodontology*. 2013; 40(5):457-73. <https://doi.org/10.1111/jcpe.12053>
 25. Fujiwara N, Tsuruda K, Iwamoto Y, Kato F, Odaki T, Yamane N, Hori Y, Harashima Y, Sakoda A, Tagaya A, Komatsuzawa H. Significant increase of oral bacteria in the early pregnancy period in Japanese women. *Journal of investigative and clinical dentistry*. 2017; 8(1):e12189. <https://doi.org/10.1111/jicd.12189>
 26. Yuan K, Wing LY, Lin MT. Pathogenetic roles of angiogenic factors in pregnancy granulomas in pregnancy are modulated by female sex hormones. *Journal of periodontology*. 2002; 73(7):701-8. <https://doi.org/10.1902/jop.2002.73.7.701>
 27. Yuan K, Lin MT. The roles of vascular endothelial growth factor and angiopoietin-2 in the regression of pregnancy pregnancy tumor. *Oral diseases*. 2004; 10(3):179-85. <https://doi.org/10.1046/j.1601-0825.2003.00997.x>
 28. Cardoso JA, Spanemberg JC, Cherubini K, FIGUEIREDO MA, Salum FG. Oral tumor gravidarum: a retrospective study of 41 cases in Southern Brazil. *Journal of Applied Oral Science*. 2013; 21(3):215-8. <https://doi.org/10.1590/1679-775720130001>
 29. Chamani G, Navabi N, Abdollahzadeh S. Prevalence of pregnancy tumor in pregnant women. *Journal of Dentistry*. 2009; 10(1):79-82.
 30. Pirie M, Cooke I, Linden G, Irwin C. Dental manifestations of pregnancy. *The Obstetrician & Gynaecologist*. 2007; 9(1):21-6. <https://doi.org/10.1576/toag.9.1.021.27292>
 31. AL-Sultani HF. Prevalence and Severity of Dental Caries, Periodontal Diseases and Dental Erosion among (20-40) Years Old Pregnant Women in Hilla city, Babylon governorate-Iraq. *Medical Journal of Babylon*. 2013; 10(2):413-20.
 32. Sarifakioglu E, Gunduz C, Gorpelioglu C. Oral mucosa manifestations in 100 pregnant versus non-pregnant patients: an epidemiological observational study. *European Journal of Dermatology*. 2006; 16(6):674-6.