

CaseReport: A Method of Impression-making for Massive Maxillectomy Defects



Fatemeh Salehi¹, Farbod Manshaee^{2*}, Majid Abolhasani²

¹ Department of Prosthodontics, School of Dentistry, Shahid Beheshti University of Medical Sciences, Tehran, Iran

² Department of Prosthodontics, Dental Implants Research Center, Dental Research Institute, School of Dentistry, University of Medical Sciences, Isfahan, Iran



Citation Salehi F, Manshaee F, Abolhasani M. A Method of Impression-making for Massive Maxillectomy Defects. Journal of Dentomaxillofacial Radiology, Pathology and Surgery. 2023; 12(4):1-4.

 <http://dx.doi.org/10.32592/3dj.12.4.1>

ABSTRACT



Article info:

Received: 22 Jul 2023

Accepted: 09 Sep 2023

Available Online: 27 Sep 2023

Maxillary defects occur for various reasons, such as congenital malformations, cancer, pathological changes, radiation burns or surgical interventions. Patients with large defects, progressive disease and those undergoing post-operative chemotherapy or radiotherapy must be rehabilitated by obturator prostheses. Making a well-extended impression for obturator prostheses may be difficult in the case of massive defects. This article presents a method for making impressions of massive maxillary defects using a special tray with a wide opening in the palatal region.

Keywords:

Neoplasms
Obturator
Palatal

* Corresponding Authors:

Farbod Manshaee.

Address: Department of Prosthodontics, Dental Implants Research Center, Dental Research Institute, School of Dentistry, Isfahan University of Medical Sciences, Isfahan, Iran

Tel: +989132265168

E-mail: fmanshaee@gmail.com

1. Introduction

Maxillary defects occur for various reasons, such as congenital malformations, cancer, pathological changes, radiation burns or surgical interventions (1).

Maxillary defects can lead to nasalized speech, dysphasia, liquid permeation to nasal cavity, loss of support of the cheek and lip, compromised esthetic and impaired mastication (2).

Prosthetic or surgical procedures may be implemented to treat this situation. Patients with large defects, progressive disease and patients undergoing post-operative chemotherapy or radiotherapy must be rehabilitated by obturator prostheses.

An obturator prosthesis is defined as "a maxillofacial prosthesis used to close a congenital or acquired tissue opening, primarily of the hard palate and/or contiguous alveolar or soft tissue structures" (3,4,5).

The vertical segment of the prosthesis provides retention, stability, oronasal separation and improves speech (6).

Accurate and properly-extended impression of areas correlated to the vertical segment is necessary for achieving satisfactory results. Meanwhile, making a well-extended impression may be difficult in the case of massive defects. This article introduces a technique for making impressions of obturator prostheses in large maxillary defects.

2. Case Report

A patient with massive maxillary defect was referred to a private dental office to receive obturator prosthesis. Premaxilla, alveolar ridges and hard palate were fully resected with the exception of the left third molar and surrounding alveolar bone (Figure 1). The primary impression was obtained using irreversible hydrocolloid (Blueprint Dentsply Sirona, Germany). The lesion boundaries were marked on the study cast (Figure 1). The special tray was fabricated from autopolymerizing acrylic resin (Special Tray Material, Dentsply). A wide opening was prepared in the palatal region of the tray, in front of the center of the maxillary defect (Figure 2). High-fusing compound (Kerr Impression Compound) was used in the first stage of the border molding procedure. For the bulb area, the compound was applied on the tray tissue surface and the tray was carried into the mouth; then, the index finger was used through the tray opening to push the

softened material into the surrounding structures: The bony undercuts of maxillary sinuses laterally, nasal floor anteriorly, and pharyngeal surface of the soft palate posteriorly (Figures 3A and 3B). The procedure was performed step by step to fully capture the bulb area and retentive structures. Compound extensions were assessed in each step and the necessary relief was taken to prevent locking of the tray in the undercuts. After completion of the border molding, the high-fusing compound was reduced by 1 mm on the surface and relined by low-fusing compound (Iso Functional Compound, GC America) in the second step (Figure 4A). The tray opening was closed by the high-fusing compound for testing the seal before taking the final impression (Figure 4B). The final impression was made with light-bodied condensation silicone (Speedex, Coltene, Switzerland) (Figure 4C). The final prosthesis is shown in Figure 5.



Figure 1. Study cast. The lesion borders are marked



Figure 2. Tray and central wide opening



Figure 3: Border molding. 3a.Finger access through the opening, 3b.Compound is extended into the lesion by finger




Figure 4: Border Molding. 4a. Finalized border molding, 4b. The tray opening closed by the compound, 4c.Final impression



Figure 5 Final prosthesis

3. Discussion

When the maxillary defect extends from the nasal cavity to the maxillary sinus, taking an exact impression can be difficult depending on the shape, size, position, and surrounding tissues.

An obturator prosthesis impression should provide proper extension into the internal structures of the

maxillary defect to provide optimum retention, stability and support. Carrying the impression compound into the lesion to properly capture the undercuts may be difficult and time-consuming. The opening in the palatal area of the tray provides 360-degree access to the internal structures of the defect and enables carrying the compound directly by the finger, thus making the procedure easy and fast and facilitating optimum

extensions.

In 2011, Bhasin et al. treated a patient with a large maxillary defect using a closed, hollow bulb obturator to improve the retention and stability of the obturator. Their applied mechanism consisted of reducing the cantilever forces of the suspension and preventing overloading of the remaining support structures (7).

Contrary to this study, in 2017, Parameswari et al. described two maxillectomy patients with large defects who were rehabilitated by a two-piece hollow bulb obturator that was retained with magnets. They suggested that rehabilitation by fabricating two-piece obturators provides light-weight prosthesis that can be easy to use, as its segments involve the undercut that improves retention (8).

In 2023, Colvenkar et al. treated a patient with palatal cleft by fabricating complete denture with obturator bulb. They also emphasized the importance of a good extension and exact impression for achieving proper retention to improve the patient's quality of life (9).

4. Conclusion

Maxillofacial prosthetics can strongly improve quality of life in patients with maxillectomy.

Given the results of this report, it can be argued that the use of this method can enable easy, fast and optimum impression-making for massive maxillary lesions.

References

1. Keyf F. Obturator prostheses for hemimaxillectomy patients. *J Oral Rehabil.* 2001;28(9):821-9. [DOI: [10.1046/j.1365-2842.2001.00754.x](https://doi.org/10.1046/j.1365-2842.2001.00754.x)] [PMID]
2. Sumita YI, Hattori M, Namba T, Ino S. Obturators to facilitate speech and swallowing in a maxillectomy patient with dementia and cerebral infarction. *Int J Maxillofac Prosthet.* 2019; 2(1): 33-5. [DOI: [10.26629/ijmp.2019.07](https://doi.org/10.26629/ijmp.2019.07)]
3. Ferro KJ, Morgano SM, Driscoll CF, Freilich MA, Guckes AD, Knoernschild KL. Glossary of prosthodontic terms: ninth edition. *J Prosthet Dent.* 2017; 117(5S): e1-e105. [DOI: [10.1016/j.prosdent.2016.12.001](https://doi.org/10.1016/j.prosdent.2016.12.001)] [PMID]
4. Phasuk K, Haug SP. Maxillofacial prosthetics. *Oral Maxillofac. Surg Clin N Am.* 2018; 30(4):487-97. [DOI: [10.1016/j.coms.2018.06.009](https://doi.org/10.1016/j.coms.2018.06.009)] [PMID]
5. Rosen EB, Palin CL, Huryn JM, Wong RJ. The role of maxillofacial prosthetics for the surgically treated patient at National Cancer Institute-designated comprehensive cancer centers. *Laryngoscope.* 2019; 129(2): 409-14. [DOI: [10.1002/lary.27330](https://doi.org/10.1002/lary.27330)] [PMID] [PMCID]
6. Kwon HB, Chang SW, Lee SH. The effect of obturator bulb height on speech in maxillectomy patients. *J Oral Rehabil.* 2011;38(3):185-95. [DOI: [10.1111/j.1365-2842.2010.02139.x](https://doi.org/10.1111/j.1365-2842.2010.02139.x)] [PMID]
7. Bhasin AS, Singh V, Mantri SS. Rehabilitation of patient with acquired maxillary defect, using a closed hollow bulb obturator. *Indian J Palliat Care.* 2011; 17(1): 70-3. [DOI: [10.4103/0973-1075.78453](https://doi.org/10.4103/0973-1075.78453)] [PMID] [PMCID]
8. Parameswari D, Rajakumar M, Jagadesaan N, Annapoorni H. Case presentation of two maxillectomy patients restored with two-piece hollow bulb obturator retained using two different types of magnets. *J Pharm Bioallied Sci.* 2017; 9(1): S252-6. [DOI: [10.4103/jpbs.JPBS_85_17](https://doi.org/10.4103/jpbs.JPBS_85_17)] [PMID] [PMCID]
9. Colvenkar S, Prakash R, Fatima S, Ahmed MS, Reddy GK. Prosthodontic Rehabilitation of a Completely Edentulous Patient With a Cleft Palate: A Case Report. *Cureus.* 2023; 15(1): e33522. [DOI: [10.7759/cureus.33522](https://doi.org/10.7759/cureus.33522)] [PMID] [PMCID].

Nevertheless, it is not possible to state the exact preferred method based on the results of a single case report, and future studies with larger sample sizes and longer follow-up periods are required.

Ethical Considerations

Compliance with ethical guidelines

Funding

None.

Authors' contributions

Fatemeh Salehi: Conceptualization, Methodology, Writing - Review & Editing Farbod Manshaee: Writing - Original Draft, Data Curation, Supervision Majid Abolhasani: Resources, Investigation, Visualization.

Conflict of Interests

The authors declare no conflict of interest.

Availability of data and material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

None.