

Research Paper: Oral Squamous Cell Carcinoma in Zahedan, Southeast Iran (2021–2023): A Cross-sectional Study



Amir Reza Khazaei^{1*}, Amir Masoud Khazaei², Alireza Gholami³, Sirus Risbaf⁴, Khashayar Moravej⁵

1. Faculty of Dentistry, Zahedan University of Medical Sciences, Zahedan, Iran.
2. Faculty of Dentistry, Mashhad University of Medical Sciences, Mashhad, Iran.
3. Student Research Committee, Faculty of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran
4. Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Zahedan University of Medical Sciences, Zahedan, Iran.
5. School of Dentistry, Islamic Azad University, Isfahan Branch, Isfahan, Iran



Citation Khazaei AR, Khazaei AM, Gholami A, Risbaf S, Moravej Kh. Oral Squamous Cell Carcinoma in Zahedan, Southeast Iran (2021–2023): A Cross-sectional Study. Journal of Dentomaxillofacial Radiology, Pathology and Surgery. 2025; 14(3): 15-20



<http://dx.doi.org/10.32592/3dj.14.2.->

ABSTRACT



Article info:

Received: 10 July 2025

Accepted: 12 Aug 2025

Available Online: 21 Aug 2025

Keywords:

- * Epidemiology
- * Mouth Neoplasms
- * Tobacco Use Cessation

Introduction: Oral squamous cell carcinoma (OSCC) is the most common malignancy of the oral cavity and a persistent public health concern. Although data exist from several regions of Iran, epidemiological evidence from the Southeast, particularly Zahedan, is scarce.

Materials and Methods: This cross-sectional study included all histopathologically confirmed OSCC cases diagnosed at the Zahedan School of Dentistry between January 2021 and December 2023 (n = 63). Demographic, clinical, and behavioral information (cigarette, hookah, smokeless tobacco/snuff, and opium use) was collected from clinical records and structured interviews. Data were analyzed statistically ($\alpha = 0.05$).

Results: The mean age was 54.2 ± 12.9 years (range 36–85), and 54.0% were women. Most participants (81.0%) self-identified as Baluch, reflecting the catchment population. The floor of the mouth was the most frequent site (38.1%), followed by mandibular gingiva (28.6%) and tongue (12.7%). Reported exposures included snuff (67.3%), cigarettes (54.0%), hookah (54.0%), and opium (47.6%); categories were not mutually exclusive. At presentation, 79.4% of lesions measured 3 cm or larger. Delays in seeking care beyond three months were more common among patients aged ≥ 50 years ($P = 0.012$) and smokers ($P = 0.019$). Pain as a presenting symptom was reported more often by women ($P = 0.020$) and patients under 50 years ($P = 0.023$).

Conclusions: In Southeast Iran, OSCC frequently presented with large lesions, floor-of-mouth predominance, and high rates of self-reported smokeless tobacco use. These findings underscore the need for early detection, culturally tailored cessation programs, and public education.

1. Introduction

Oral squamous cell carcinoma (OSCC), which is over 90% of the oral malignancy, is continuing to be a significant global health challenge (1). It is the 11th most frequent cancer in men and the 16th in women, representing about 27,000 new cases and 5,500 deaths each year (2). Well-established risk factors include tobacco in various forms—such as betel quid, pan masala, gutka, and smokeless tobacco—as well as cigarette smoking, alcohol, viral infections, and immunosuppression (3, 4). In many Asian regions, particularly India and Taiwan, the high prevalence of smokeless tobacco use has been directly linked to

increased OSCC burden (3).

Clinical presentation of OSCC is variable, ranging from leukoplakia and erythroplakia to ulcerated or exophytic masses with irregular borders (5-7). Common intraoral sites at risk include the lower lip, floor of mouth, and lateral tongue borders (3). Although histopathological confirmation through biopsy is essential for diagnosis (8), patients frequently present at advanced stages when treatment requires multimodal approaches and prognosis is poor (6, 9-11).

Although some national studies have been conducted, there is a paucity of epidemiological studies in Southeast

* Corresponding Authors:

Amir Reza Khazaei

Address: Faculty of Dentistry, Zahedan University of Medical Sciences, Zahedan, Iran

E-mail: khazaeiamirreza@outlook.com

Iran. Zahedan, the capital city of Sistan and Baluchestan Province, is a deprived area with high prevalence of smokeless tobacco (SLT) and opium consumption and with possible poor access to health care. It is important to describe the demography and behavioural profile of OSCC patients in this setting in order to plan for appropriate screening and preventive strategies.

This study aimed to characterize demographic, clinical and behavioral characteristics in cases of OSCC with histopathologically-confirmed diagnosis that were diagnosed between 2021 and 2023 in Zahedan, and to investigate simple associations of the variables (age, sex and reported habits) with the clinical profile.

2. Materials and Methods

This is a cross-sectional study that was conducted between 2021 and 2023. The study population was patients with oral squamous cell carcinoma referred to the Dentistry School of Zahedan, Iran. Ethics approval was obtained from the University of Zahedan Research Ethics Board [IR.ZAUMS.REC.1400.100].

In this study, 63 patients with a confirmed OSCC who referred to Zahedan Dental School from 2021 to 2023, were investigated as a census. The inclusion criteria were all patients with histopathologically confirmed primary OSCC diagnosed between 2021 and 2023, and the exclusion criteria were non-OSCC malignancies, recurrent or previously treated OSCC, cases without histological confirmation, and patients with incomplete demographic or clinical data.

After patients suspected of oral malignant lesions came to the Dental School of Zahedan, the clinical examinations were performed by Oral Medicine Specialists or Oral and Maxillofacial surgeons. Also, a definite diagnosis was made by Histological analyses at the Oral and Maxillofacial Pathology Department of the Zahedan Dentistry School.

Once patients were diagnosed with OSCC and accepted the conditions of the research, a senior dentistry student under the supervision of oral medicine specialists collected data through clinical oral examinations and a questionnaire consisting of demographics, behavioral risk factors, involved intraoral regions and the largest dimension of the lesion, and symptoms which patients were reporting at the time of clinical examination, and duration of lesion presence in mouth since the first time, the patient was

aware of its appearance.

Demographic, clinical, and behavioral information (cigarette, hookah, smokeless tobacco/snuff, and opium use) were obtained from clinical records and structured interviews. Behavioral variables were recorded as yes/no. Although frequency and duration of use were not systematically assessed, all participants were asked standardized questions. Face and content validity of the questionnaire were reviewed by oral medicine specialists, and reliability was evaluated in a pilot sample (Cronbach's $\alpha = 0.82$).

Statistical analyses were performed using SPSS software (version 22, IBM Corp., Armonk, NY, USA). Descriptive statistics were calculated as means $\pm$ standard deviation for continuous variables and as frequencies and percentages for categorical variables. Associations between categorical variables such as age group, gender, tobacco and snuff use, lesion site, and delay in seeking treatment were assessed using the Chi-square test or Fisher's exact test when expected cell counts were small. Normality of continuous variables was checked using the Shapiro-Wilk test. All statistical tests were two-tailed, and a significance threshold of $P < 0.05$ was applied. Subgroup analyses by gender and age were considered exploratory.

3. Results

Sixty-three patients with histopathologically confirmed oral squamous cell carcinoma (OSCC) were included. The mean age was 54.2 ± 12.9 years (range, 36–85), and 55.6% of the participants were older than 50 years. Women comprised 54.0% of cases ($n = 34$). A family history of oral cancer was reported by 31.7% of patients, and 81.0% ($n = 51$) self-identified as Baluch, reflecting the local catchment population (Table 1).

The most common habit was snuff (naswar) use in 42 patients (66.7%), followed by cigarette smoking in 34 (54.0%), hookah in 34 (54.0%), and opium in 30 (47.6%) (Figure 1).

At presentation, 34.9% of patients reported pain only, 15.9% reported pain with burning, 33.3% reported pain combined with burning and hemorrhage, and 15.9% were asymptomatic. The interval from first awareness of the lesion to seeking care was 1–3 months in 41.3% and 3–6 months in 42.9% of patients. Most lesions (52.4%) measured 3–10 cm; overall, 79.4% were larger than 3 cm (Table 2).

Table 1. Demographic characteristics of the patients with oral squamous cell carcinoma in Zahedan, southeast of Iran, 2021-2023

		Cases	Percent	Total
Gender	Male	29	46.0%	63 (100%)
	Female	34	54.0%	
Race	Fars	12	19.0%	63 (100%)
	Baluch	51	81.0%	
Family history of oral cancer	Yes	20	31.7%	63 (100%)
	No	43	68.3%	
Age (Mean $\pm$ SD, Range)		54.24 $\pm$ 12.9, 36-85		

Figure 1. Distribution of some of the most common risk factors among patients with Oral Squamous Cell Carcinoma in Zahedan, Southeast of

Iran, 2021-2023




Table 2. Distribution of symptoms, Size of the lesion, and duration of the lesion's presence among patients with oral squamous cell carcinoma in Zahedan, southeast of Iran, 2021-2023

		Cases	Percent	Total
Symptoms at the time of examination	Pain	22	34.9%	63 (100%)
	Pain and burning sensation	10	15.9%	
	Pain, burning sensation, and hemorrhage	21	33.3%	
	Without symptoms	10	15.9%	
Duration of the presence of the lesion in the mouth before searching for treatment	Less than 1month	7	11.1%	63 (100%)
	1-3 months	26	41.3%	
	3-6 months	27	42.9%	
	6-12 months	3	4.8%	
Size of the lesion	≤ 2	13	20.6%	63 (100%)
	3-10	33	52.4%	
	> 10	17	27.0%	

The most frequent primary site was the floor of the mouth (38.1%), followed by mandibular gingiva (28.6%) and tongue (12.7%). Buccal mucosa was the least common (1.6 %; [Figure 2](#)).

Longer delays in presentation (> 3 months) were more common in patients aged ≥ 50 years compared with younger patients (57.1% vs. 25.0%; $P = 0.012$) and in smokers compared with non-smokers (55.6% vs. 11.1%; $P = 0.019$; [Table 3](#)). For instance, lesions present for 3–6 months were reported in 57.1% of older patients compared

with 25.0% of younger patients, and in 55.6% of smokers compared with 11.1% of non-smokers.

Pain as a presenting symptom was more frequent in women than men (44.1% vs. 24.1%; $P = 0.020$) and in patients younger than 50 years compared with older patients (42.9% vs. 28.6%; $P = 0.023$) ([Table 4](#)). No other significant associations were observed between demographic or behavioral factors and tumor site, size, or symptoms.

Figure 2. Distribution of intraoral location of the lesions in patients with oral squamous cell carcinoma in Zahedan, southeast of Iran, 2021-2023




Table 3. Distribution of Duration of the presence of the lesion in the mouth before searching for treatment among patients with oral squamous cell carcinoma according to smoking and age in southeast of Iran.

Duration of the presence of the lesion in the mouth before searching for treatment			< 1 month	1-3 months	3-6 months	6-12 months	Total	P-value
Age	≤ 50	N	4	16	7	1	28	0.012
		%	14.2%	57.2%	25%	3.6%	100.0%	
	> 50	N	3	10	20	2	35	
		%	8.6%	28.5%	57.2%	5.7%	100.0%	
Smoking	Yes	N	2	16	25	2	45	0.019
		%	4.5%	34.5%	56.0%	4.5%	100.0%	
	No	N	5	10	2	1	18	
		%	27.8%	55.5%	11.2%	5.5%	100.0%	



Test: Chi-square or Fisher's exact, as appropriate. Significance: $\alpha = 0.05$.

Table 4. Distribution of symptoms among oral squamous cell carcinoma patients by age and smoking status.

Symptoms at the time of examination			Pain	Pain and burning sensation	Pain, burning sensation, and hemorrhage	Without symptoms	Total	P-value
Sex	Male	N	7	9	9	4	29	0.020
		%	24.1%	31.0%	31.0%	13.8%	100.0%	
	Female	N	15	1	12	6	34	
		%	44.1%	2.9%	35.3%	17.6%	100.0%	
Age (year)	≤ 50	N	12	0	11	5	28	0.023
		%	42.9%	0.0%	39.3%	17.9%	100.0%	
	> 50	N	10	10	10	5	35	
		%	28.6%	28.6%	28.6%	14.3%	100.0%	



Test: Chi-square or Fisher's exact, as appropriate. Significance: $\alpha = 0.05$.

4. Discussion

This cross-sectional study provides updated information on the epidemiology of oral squamous cell carcinoma (OSCC) in Southeast Iran. The mean age of patients was 54.2 years, which is lower than that reported in some Brazilian series (12, 13) but higher than findings from Canada (14). Women represented 54% of cases, a proportion differing from several reports where men predominated (12, 14, 15). However, studies from South

Asia have also reported higher prevalence among women (16), possibly reflecting cultural practices that discourage smoking among women but allow smokeless tobacco use. Such explanations should be interpreted cautiously given our descriptive design.

Snuff use was the most frequently reported risk factor (67.3%), consistent with studies demonstrating the role of smokeless tobacco in oral cancer (15-18). In our series, the floor of the mouth was the most common tumor site

(38.1%), aligning with some prior findings (19, 20), although other populations have reported the tongue or lip as more common locations.

Delayed presentation was frequent: 42.9% of patients, mainly older individuals and smokers, sought treatment only after 3–6 months. Such delays highlight the lack of awareness regarding early signs of OSCC and the absence of systematic screening programs in Iran. Pain as a presenting symptom was more often reported by younger patients and women, which may reflect differences in health-seeking behavior or symptom perception rather than biological variation. The predominance of Baluch patients (81.0%) in this cohort likely reflects the regional catchment population rather than ethnicity as an independent risk factor. Future analytic studies, such as case–control designs, are required to clarify potential associations.

The present findings highlight the importance of early detection and culturally sensitive risk-reduction strategies in regions with high prevalence of smokeless tobacco use. For clinicians, careful inspection of high-risk sites such as the floor of the mouth and mandibular gingiva is warranted, even in relatively young patients or those presenting with non-specific symptoms such as burning sensation. Integrating routine oral cancer screening into dental visits, combined with brief chairside counseling on naswar and other risk habits, could reduce diagnostic delays and ultimately improve prognosis.

Strengths of the present study include complete case ascertainment during the study period and pathology-confirmed diagnoses. Limitations include the single-center design, modest sample size, reliance on self-reported behaviors (subject to recall and social desirability bias), lack of detailed exposure frequency/duration, and absence of multivariable analysis.

These findings emphasize the need for public health programs promoting awareness of OSCC risk factors, particularly smokeless tobacco, and the benefits of early diagnosis. Establishing oral cancer screening in dental clinics and integrating culturally tailored cessation programs into community health strategies could help reduce diagnostic delays and disease burden in Southeast Iran (10).

References

1. Badwelan M, Muaddi H, Ahmed A, Lee KT, Tran SD. Oral squamous cell carcinoma and concomitant primary tumors, what do we know? A review of the literature. *Curr Oncol*. 2023;30(4):3721-34. [DOI: 10.3390/currenco30040283] [PMID] [PMCID]
2. Neville BW, Damm DD, Allen CM, Chi AC. *Oral and maxillofacial pathology*. Elsevier Health Sciences; 2015. [Link]
3. Glick M, Greenberg MS, Lockhart PB, Challacombe SJ. Introduction to oral medicine and oral diagnosis: patient evaluation. *Burket's Oral Medicine*. 2021;1-18. [Link]
4. Dalirsani Z, Ghazi A, Pakfetrat A, Jamali J, Foroughi Z, Mirzaei M. The relationship between smokeless tobacco and the incidence of oral cancer: a systematic review study. *Addict Health*. 2025;17(1):1-15. [DOI: 10.34172/ahj.1522] [PMID] [PMCID]
5. Kawczak P, Feszak I, Bączek T. Comparative mechanistic insights and therapeutic potential of pembrolizumab, durvalumab, and ipilimumab in the targeted management of oral and head and neck squamous cell carcinoma. *Cancers*. 2025;17:2805. [DOI: 10.3390/cancers17172805] [PMID] [PMCID]
6. Kijowska J, Grzegorzczak J, Gliwa K, Jędras A, Sitarz M. Epidemiology, diagnostics, and therapy of oral cancer—update review. *Cancers*. 2024;16(18):3156. [DOI: 10.3390/cancers16183156] [PMID] [PMCID]
7. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*.

5. Conclusions

In this cross-sectional study from Southeast Iran, most OSCC cases presented at relatively advanced stages with large lesions and frequent involvement of the floor of the mouth. High rates of self-reported smokeless tobacco use were observed, alongside substantial delays in seeking care, particularly among older patients and smokers. While causality cannot be inferred, these findings highlight the need for public awareness programs, culturally tailored tobacco cessation initiatives, and improved referral systems to facilitate earlier diagnosis and management of OSCC in underserved regions.

Ethical Considerations

This study was approved by the Ethics Committee of Zahedan University of Medical Sciences (Approval No. IR.ZAUMS.REC.1400.100).

Funding

The authors thank Zahedan University of Medical Sciences for their approval to conduct this project.

Authors' Contributions

Amir Reza Khazaei: Conceptualization; Supervision; Formal Analysis **Amir Masoud Khazaei:** Investigation, Writing – Original Draft **Alireza Gholami:** Resources, Data Curation **Sirus Risbaf:** Project Administration, Writing – Review & Editing **Khashayar Moravej:** Methodology, Writing – Review & Editing.

Conflict of Interests

The authors declare no conflict of interest.

Availability of Data and Material

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

None.

- 2024;74(3):229-63. [DOI: [10.3322/caac.21834](https://doi.org/10.3322/caac.21834)] [PMID]
8. Wong T, Wiesenfeld D. Oral cancer. *Aust Dent J*. 2018;63(S1):S91-S99. [DOI: [10.1111/adj.12594](https://doi.org/10.1111/adj.12594)] [PMID]
9. Mahmood BJ, Gataa IS, Mustafa UM. Clinicopathological presentation of oral squamous cell carcinoma in Iraqi population. *J Craniomaxillofac Res*. 2021;8(3):129-142. [Link]
10. Rumgay H, Nathan ST, Shah R, Vignat J, Ayo-Yusuf O, Chaturvedi P, et al. Global burden of oral cancer in 2022 attributable to smokeless tobacco and areca nut consumption: a population attributable fraction analysis. *Lancet Oncol*. 2024;25(11):1413-23. [DOI: [10.1016/s1470-2045\(24\)00458-3](https://doi.org/10.1016/s1470-2045(24)00458-3)] [PMID]
11. Katirachi SK, Grønlund MP, Jakobsen KK, Grønhoj C, von Buchwald C. The prevalence of HPV in oral cavity squamous cell carcinoma. *Viruses*. 2023;15(2):451. [DOI: [10.3390/v15020451](https://doi.org/10.3390/v15020451)] [PMID] [PMCID]
12. Louredo BVR, Vargas PA, Pérez-de-Oliveira ME, Lopes MA, Kowalski LP, Curado MP. Epidemiology and survival outcomes of lip, oral cavity, and oropharyngeal squamous cell carcinoma in a southeast Brazilian population. *Med Oral Patol Oral Cir Bucal*. 2022;27(3):e274. [DOI: [10.4317/medoral.25147](https://doi.org/10.4317/medoral.25147)] [PMID] [PMCID]
13. Solhi M, Fattahi E, Barati H, Mohammadi M, Kasmaei P, Rastaghi S. Smokeless tobacco use in Iran: a systematic review. *Addict Health*. 2020;12(3):225-34. [DOI: [10.22122/ahj.v12i3.274](https://doi.org/10.22122/ahj.v12i3.274)] [PMID] [PMCID]
14. Badri P, Lai H, Ganatra S, Baracos V, Amin M. Factors associated with oral cancerous and precancerous lesions in an underserved community: a cross-sectional study. *Int J Environ Res Public Health*. 2022;19(3):1297. [DOI: [10.3390/ijerph19031297](https://doi.org/10.3390/ijerph19031297)] [PMID] [PMCID]
15. Colares N, Souza Rodrigues DF, Freitas MO, Dantas TS, Cunha M, Sousa FB, et al. Smoking history decreases survival in patients with squamous cell carcinoma of the mouth: a retrospective study with 15 years of follow-up. *Asian Pac J Cancer Prev*. 2019;20(6):1781-7. [DOI: [10.31557/apjcp.2019.20.6.1781](https://doi.org/10.31557/apjcp.2019.20.6.1781)] [PMID] [PMCID]
16. Toralla O, Lopez Jornet P, Pons-Fuster E. The effect of an informative video upon anxiety and stress in patients requiring an oral biopsy: a randomized controlled study. *Int J Environ Res Public Health*. 2022;19(2):783. [DOI: [10.3390/ijerph19020783](https://doi.org/10.3390/ijerph19020783)] [PMID] [PMCID]
17. Graboyes EM, Kompelli AR, Neskey DM, Brennan E, Nguyen S, Sterba KR, et al. Association of treatment delays with survival for patients with head and neck cancer: a systematic review. *JAMA Otolaryngol Head Neck Surg*. 2019;145(2):166-77. [DOI: [10.1001/jamaoto.2018.2716](https://doi.org/10.1001/jamaoto.2018.2716)] [PMID] [PMCID]
18. da Cunha Lima AM, Meira IA, Soares MSM, Bonan PRF, Melo CB, Dalle Piagge CSL. Delay in diagnosis of oral cancer: a systematic review. *Med Oral Patol Oral Cir Bucal*. 2021;26(6):e815-e824. [DOI: [10.4317/medoral.24808](https://doi.org/10.4317/medoral.24808)] [PMID] [PMCID]
19. Anwar N, Chundrager Q, Awan S, Moatter T, Ali TS, Abdul Rasheed M, et al. Prevalence of high-risk human papillomavirus in oral squamous cell carcinoma with or without chewing habits. *PLoS One*. 2024;19(5):e0300354. [DOI: [10.1371/journal.pone.0300354](https://doi.org/10.1371/journal.pone.0300354)] [PMID] [PMCID]
20. Ha PK, Califano JA. The role of human papillomavirus in oral carcinogenesis. *Crit Rev Oral Biol Med*. 2004;15(4):188-96. [DOI: [10.1177/154411130401500402](https://doi.org/10.1177/154411130401500402)] [PMID]